

EXTENSIONS OF THE REPRESENTATION MODULES OF A PRIME ORDER GROUP

V. A. BOVDI, V. P. RUDKO

Dedicated to Professor Kálmán Győry on his 65th birthday

ABSTRACT. For the ring R of integers of a ramified extension of the field of p -adic numbers and a cyclic group G of prime order p we study the extensions of the additive groups of R -representations modules of G by the group G .

Let $\mathfrak{F}$ be the field of fractions of a principal ideal domain R , F a field which contains R , let G be a finite group and Γ a matrix R -representation of G . Let M be an RG -module, which affords the R -representation Γ of G , and $FM = F \otimes_R M$ the smallest linear space over F which contains M and $\widehat{M} = FM^+/M$, the factor group of the additive group of the space FM by the additive group of M . Clearly, the group $\widehat{M}$ and the space FM are RG -modules. Put $\widehat{F} = F^+/R$.

Let $\mathfrak{f} : G \rightarrow \widehat{M}$ be a 1-cocycle of G with value in $\widehat{M}$, i.e.

$$\mathfrak{f}(xy) = x\mathfrak{f}(y) + \mathfrak{f}(x), \quad (x, y \in G).$$

Define $[g, x]$ by $\begin{pmatrix} g & x \\ 0 & 1 \end{pmatrix}$ and set

$$\mathfrak{Crns}(G, M, \mathfrak{f}) = \{ [g, x] \mid g \in G, \quad x \in \mathfrak{f}(g) \},$$

where x runs over the cosets $\mathfrak{f}(g) \in \widehat{M}$ for any $g \in G$.

Clearly, $\mathfrak{Crns}(G, M, \mathfrak{f})$ is a group, where the multiplication is the usual matrix multiplication. Of course $K_1 = \{[e, x] \mid e \text{ is the unit element of } G, \quad x \in \mathfrak{f}(e)\}$ is a normal subgroup of $\mathfrak{Crns}(G, M, \mathfrak{f})$ such that $K_1 \cong M^+$ and $\mathfrak{Crns}(G, M, \mathfrak{f})/K_1 \cong G$. The group $\mathfrak{Crns}(G, M, \mathfrak{f})$ is an extension of the additive group of the RG -module M by G .

We are using the terminology of the theory of group representations [1].

A 1-cocycle $\mathfrak{f} : G \rightarrow \widehat{M}$ is called *coboundary*, if there exists an $x \in FM$ such that $\mathfrak{f}(g) = (g - 1)x + M$ for every $g \in G$. The 1-cocycles $\mathfrak{f}_1 : G \rightarrow \widehat{M}$ and

1991 *Mathematics Subject Classification*. Primary 16U60, 16W10.

The research was supported by OTKA No.T 037202 and No.T 038059

$f_2 : G \rightarrow \widehat{M}$ are called *cohomologous* if $f_1 - f_2$ is a coboundary. Let $H^1(G, \widehat{M})$ be the first cohomology group. Clearly, each element of $H^1(G, \widehat{M})$ defines a class of equivalence of groups.

If the 1-cocycles f_1, f_2 are cohomologous, then $\mathfrak{C}r\eta\mathfrak{s}(G, M, f_1)$ and $\mathfrak{C}r\eta\mathfrak{s}(G, M, f_2)$ are isomorphic. This isomorphism is called *equivalence* and these groups are called equivalent. In particular, the group $\mathfrak{C}r\eta\mathfrak{s}(G, M, f)$ is split (i.e. $\mathfrak{C}r\eta\mathfrak{s}(G, M, f) = M \rtimes G$) if and only if f is coboundary.

The *dimension* of the group $\mathfrak{C}r\eta\mathfrak{s}(G, M, f)$ is called the R -rank of the R -module M . (Note that M is a free R -module of finite rank.) The group $\mathfrak{C}r\eta\mathfrak{s}(G, M, f)$ is called *irreducible* (*indecomposable*), if M is an irreducible (indecomposable) RG -module and the 1-cocycle f is not cohomologous to zero.

The group $\mathfrak{C}r\eta\mathfrak{s}(G, M, f)$ is *non-split*, if the 1-cocycle f defines a nonzero element of $H^1(G, \widehat{M})$.

Note that the properties of the group $\mathfrak{C}r\eta\mathfrak{s}(G, M, f)$ were studied in [5,6,8], in the cases when R is either the ring of rational integers $\mathbb{Z}$, or the p -adic integers $\mathbb{Z}_p$, or the localization $\mathbb{Z}_{(p)}$ of $\mathbb{Z}$ at p .

Let $G = \langle a \mid a^p = 1 \rangle$ be the cyclic group of prime order p , R the ring of integers of the ramified finite extension T of the field of p -adic numbers. We calculate the group $H^1(G, \widehat{M})$ for some module M of an indecomposable R -representation of G .

Let $\Phi_p(x) = x^{p-1} + \dots + x + 1$ be a cyclotomic polynomial of degree p and let $\eta(x)$ be a divisor of $\Phi_p(x)$ over the field $\mathfrak{T}$ with $\deg(\eta(x)) < p - 1$ (provided that such nontrivial polynomial exists).

Lemma 1. *Let M_1 and M_2 be RG -modules which afford an R -representation Γ of $G = \langle a \mid a^p = 1 \rangle$.*

(i) *If $M_1 \cong M_2$ then $H^1(G, \widehat{M}_1) \cong H^1(G, \widehat{M}_2)$.*

(ii) *If the matrix $\Gamma(a)$ does not have 1 as eigenvalue, then $H^1(G, \widehat{M}_1)$ is trivial.*

Proof. See [1]. $\square$

Theorem 1. *Let $G = \langle a \mid a^p = 1 \rangle$ and $M_\eta = \eta(a)RG$. Then the RG -module M_η is indecomposable and*

$$H^1(G, \widehat{M}_\eta) \cong R/(\eta(1)R),$$

where $R/(\eta(1)R)$ is the additive group of the factor ring of R by the ideal $\eta(1)R$.

Proof. Let $t \in R$ be a prime element and $\overline{R} = R/(tR)$. Then in $\overline{R}$ we have that

$$x^p - 1 = (x - 1)^p; \quad \eta(x) = (x - 1)^n; \quad (x^p - 1)\eta^{-1}(x) = (x - 1)^{p-n}, \quad (1)$$

where $n = \deg(\eta(x))$.

Put $\eta_1(x) = (x^p - 1)\eta^{-1}(x)$. Then M_η and $RG/(\eta_1(a)RG)$ are isomorphic as RG -modules. If $\overline{M}_\eta = M_\eta/(tM_\eta)$, then by (1) follows that $\overline{M}_\eta$ is a root subspace of the linear operator a over $\overline{R}$. It is easy to see that $\overline{M}_\eta$ is not decomposable into a direct sum of invariant subspaces. It follows that M_η is an indecomposable

RG -module. Clearly $FM_\eta = \eta(a)FG = \eta(a)F + (a-1)FM_\eta$ and the group $\widehat{M_\eta} = FM_\eta^+/M_\eta$ is isomorphic to a direct sum of groups $\eta(a)(F^+/R) + (a-1)\widehat{M_\eta}$. This means that in the class of 1-cocycles there is a cocycle $f : G \rightarrow \widehat{M_\eta}$ such that

$$f(a) = \lambda\eta(a) + M_\eta, \quad (\lambda \in F).$$

Moreover, from $f(a^p) = 0$ (in $\widehat{M_\eta}$) it follows that if $\omega = a^{p-1} + \dots + a + 1$, then

$$\omega \cdot f(a) = \lambda \cdot \eta(1)\omega \in M_\eta$$

if and only if $\lambda\eta(1) \in R$. Therefore, $H^1(G, \widehat{M_\eta})$ is isomorphic to the subgroup $\{ \lambda + R \mid \lambda \in F, \lambda \cdot \eta(1) \in R \}$ of F/R and

$$\{ \lambda + R \in F/R \mid \lambda \cdot \eta(1) \in R \} \cong R/(\eta(1)R).$$

□

Corollary 1. *Let $\alpha \in R$ and $\eta(1)R = t^s R$, where t is a prime element of R . Put*

$$K_\alpha(G, M_\eta) = \left\langle \begin{pmatrix} e & m \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} a & \alpha t^{-s} \eta(a) \\ 0 & 1 \end{pmatrix} \mid m \in M_\eta \right\rangle,$$

where α runs over the representative elements of the cosets of $R/(t^s R)$. Up to equivalence, the groups $K_\alpha(G, M_\eta)$ give all extensions of the additive group of the RG -module M_η by the group G .

Suppose $p = t^d \theta$, where $d > 1$ is the ramification index and θ is a unit in R . Set

$$\mathfrak{X}_{ji} = t^j RG + (a-1)^i RG, \quad (1 \leq j < d, 1 \leq i < p).$$

Theorem 2. *The module $\mathfrak{X}_{ji}$ is an RG -module affording an indecomposable R -representation of G and*

$$H^1(G, \widehat{\mathfrak{X}_{ji}}) \cong R/(t^{d-j} R).$$

Proof. Suppose that the RG -module $\mathfrak{X}_{ji}$ is decomposable into a direct sum of RG -submodules. Then $t^j = u_1 + u_2$, where u_1, u_2 are nonzero elements of RG with $u_1 u_2 = 0$. Thus $e_1 = t^{-j} u_1$ is an idempotent. Since the trace $tr(e_1)$ of e_1 is a rational number (see Theorem 3.5, [4] p.21) of the form rp^{-1} ($1 \leq r \leq p$), we get $t^j r p^{-1} \in R$, which is impossible for $j < d$. This contradiction proves the indecomposability of $\mathfrak{X}_{ji}$. Clearly $F\mathfrak{X}_{ji} = FG = F + (a-1)FG$. Therefore, in each class of 1-cocycles there is a cocycle $f : G \rightarrow \widehat{\mathfrak{X}_{ji}}$ such that $f(a) = \lambda + \mathfrak{X}_{ji}$, where $\lambda \in F$ with $\lambda\omega \in \mathfrak{X}_{ji}$. It follows that $\lambda p = t^j \alpha$, where $\alpha \in R$ and

$$H^1(G, \widehat{\mathfrak{X}_{ji}}) \cong \{ \lambda + R \mid \lambda \in F, \lambda t^{d-j} \in R \}$$

is a subgroup of F/R .

□

Set

$$K_\alpha(G, \mathfrak{X}_{ji}) = \left\langle \begin{pmatrix} e & m \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} a & \alpha t^{j-d} \\ 0 & 1 \end{pmatrix} \mid m \in \mathfrak{X}_{ji} \right\rangle,$$

where α runs over the representative elements of the cosets of $R/(t^{d-j} R)$.

Corollary 2. *The groups $K_\alpha(G, \mathfrak{X}_{ji})$ give all extensions of the additive group of the RG -module $\mathfrak{X}_{ji}$ by G .*

Lemma 2. *The set $\{ \mathfrak{X}_{ji} \mid j = 1, \dots, d-1; i = 1, \dots, \frac{p-1}{2} \}$ consists of pairwise non-isomorphic modules.*

Proof. Let us consider an indecomposable $\overline{RG}$ -module $V_i = \overline{RG}/((a-1)^i \overline{RG})$, where $\overline{R} = R/(tR)$ and $1 \leq i \leq p$. It is easy to check that the elements

$$u_1 = t^j, \dots, u_i = t^j(a-1)^{i-1}, \quad u_{i+1} = (a-1)^i, \dots, u_p = (a-1)^{p-1} \quad (2)$$

form an R -basis in $\mathfrak{X}_{ji}$ and

$$\Phi_p(x) - (x-1)^{p-1} = p\theta(x), \quad (3)$$

where $\theta(x) \in \mathbb{Z}[x]$, $\deg(\theta(x)) \leq p-2$. Note that since $\theta(1) = 1$, it follows that $\theta(a)$ is a unit in the group ring RG . Using the identity

$$xy - 1 = (x-1)(y-1) + (x-1) + (y-1),$$

from (3) we obtain that

$$(a-1)^p = p(a-1) \cdot (\alpha_0 + \alpha_1(a-1) + \dots + \alpha_{p-2}(a-1)^{p-2}), \quad (4)$$

where $\alpha_0, \alpha_1, \dots, \alpha_{p-2} \in \mathbb{Z}$. Since $p = t^d \theta = t(t^{d-1} \theta)$, from (4) we get

$$(a-1)^p = (a-1)u_p = tm, \quad (5)$$

where $m \in \mathfrak{X}_{ji}$. According to (2) $(a-1)u_i = t^j u_{i+1}$, and from (5) we obtain that the RG -module $\overline{\mathfrak{X}}_{ji} = \mathfrak{X}_{ji}/(t\mathfrak{X}_{ji})$ is isomorphic to a direct sum $V_i \oplus V_{p-i}$ of indecomposable $\overline{RG}$ -modules, so by Theorem 2 and Lemma 1 the proof is complete. $\square$

Let $n > 1$ be the degree of a divisor of $\Phi_p(x)$, which is irreducible over R . We consider the following RG -modules:

$$\mathfrak{U}_{ji} = t^j(a-1)RG + (a-1)^{s+1}RG, \quad (1 \leq j < d, \quad 1 \leq s < n).$$

It is easy to check that the RG -module $\mathfrak{U}_{ji}$ satisfies the condition (ii) of Lemma 1, so $H^1(G, \widehat{\mathfrak{U}}_{ji}) = 0$.

Let $\mathfrak{Z}_{js}$ be a submodule of the free module $RG^{(2)} = \{(x, y) \mid x, y \in RG\}$ of rank 2, which consists of the solutions (x, y) of the equality

$$t^j(a-1)x + (a-1)^{s+1}y = 0. \quad (6)$$

Lemma 3. *Let $\omega = \Phi_p(a)$ and set $u_1 = [0, \omega]$, $u_2 = [(a-1)^s, -t^j]$ and $u_3 = [t^{-j}(\omega - (a-1)^{p-1}), (a-1)^{p-s-1}]$. Then $\mathfrak{Z}_{js}$ is an RG -module generated by u_1, u_2, u_3 .*

Proof. Clearly, $u_1, u_2, u_3 \in \mathfrak{Z}_{js}$. Let $u = [x, y]$ be an arbitrary element of $\mathfrak{Z}_{js}$. If $x = 0$ then $u = u_1$. Suppose $x \neq 0$. By subtraction of the elements of RGu_3 from u we obtain that $y = \gamma_0 + \gamma_1(a-1) + \cdots + \gamma_{p-s-2}(a-1)^{p-s-2}$ ($\gamma_r \in R$). By (6)

$$t^j(a-1)x + (\gamma_0 + \gamma_1(a-1) + \cdots + \gamma_{p-s-2}(a-1)^{p-s-2}) \cdot (a-1)^{s+1} = 0,$$

which is possible if and only if $\gamma_0 \equiv \cdots \equiv \gamma_{p-s-2} \equiv 0 \pmod{t^j}$. Now, since u is an element of RGu_2 , we obtain that $y = 0$. Then $t^j(a-1)x = 0$ which implies $x = \alpha\omega$ ($\alpha \in R$) and $u = \alpha(t^j u_3 - (a-1)^{p-s-1} u_2)$. □

Theorem 3. *The RG -module $\mathfrak{Z}_{js}$ is indecomposable. Moreover,*

$$H^1(G, \widehat{\mathfrak{Z}}_{js}) \cong R/(t^d R) \oplus R/(t^{d-j} R)$$

and the RG -modules $\mathfrak{X}_{js}$ are pairwise non-isomorphic.

Proof. It is easy to see that

$$u_1 = t^j(a-1), \dots, u_{i-1} = t^j(a-1)^{i-1}, \quad u_i = (a-1)^i, \dots, u_{p-1} = (a-1)^{p-1}$$

form an R -basis in the RG -module $\mathfrak{U}_{js}$ and

$$\overline{\mathfrak{U}}_{js} = \mathfrak{U}_{js}/(t\mathfrak{U}_{js}) \cong V_s \oplus V_{p-s-1}.$$

Since $s < n$, it follows that the RG -module $\mathfrak{U}_{js}$ is indecomposable. Moreover, it follows that the RG -modules $\mathfrak{U}_{js}$ are pairwise non-isomorphic and RG -modules $\mathfrak{Z}_{js}$, $\mathfrak{U}_{js}$ and RG^2 form an exact sequence

$$0 \rightarrow \mathfrak{Z}_{js} \rightarrow RG^{(2)} \rightarrow \mathfrak{U}_{js} \rightarrow 0.$$

Therefore, $\mathfrak{Z}_{js}$ is the kernel of a minimal projective covering of the indecomposable RG -module $\mathfrak{U}_{js}$, so $\mathfrak{Z}_{js}$ is also indecomposable.

Lemma 4. *Let $\widehat{\mathfrak{Z}}_{js} = (F\mathfrak{Z}_{js})^+/\mathfrak{Z}_{js}$, $\widehat{F} = F^+/R$ and $M = (a-1)\widehat{\mathfrak{Z}}_{js}$. Then*

$$\widehat{\mathfrak{Z}}_{js}/M = \widehat{F}\nu_1 + \widehat{F}\nu_2,$$

where $\nu_1 = [0, \omega] + M$, $\nu_2 = [\omega, 0] + M$ and $a\nu_1 = \nu_1$, $a\nu_2 = \nu_2$.

Proof. Clearly, $ax = x$ ($x \in \widehat{\mathfrak{Z}}_{js}/M$) and $\widehat{F}\nu_1 = \widehat{F}[0, \omega] + M \in \widehat{\mathfrak{Z}}_{js}/M$. Moreover,

$$\begin{aligned} \omega\widehat{F}u_3 + M &= \widehat{F}[t^{-1}p\omega, 0] + M = \widehat{F}(tp^{-1})[t^{-1}p\omega, 0] + M \\ &= \widehat{F}[\omega, 0] + M. \end{aligned}$$

By analogy

$$\omega \widehat{F}u_2 + M = \widehat{F}[0, -t^j\omega] + M = \widehat{F}[0, \omega] = \widehat{F}\nu_1.$$

Therefore $\widehat{\mathfrak{Z}}_{js}/M = \widehat{F}\nu_1 + \widehat{F}\nu_2$. $\square$

From Lemma 4 it follows that each class of 1-cocycles of the group G with values in the group $\widehat{\mathfrak{Z}}_{js} = (F\mathfrak{Z}_{js})^+/\mathfrak{Z}_{js}$ contains a 1-cocycle $\mathfrak{f}$ such that

$$\mathfrak{f}(a) = \alpha[0, \omega] + \beta[\omega, 0] + \mathfrak{Z}_{js},$$

where $\alpha, \beta \in F$ and $\omega(\alpha[0, \omega] + \beta[\omega, 0]) \in \mathfrak{Z}_{js}$. This condition holds if and only if $\alpha p, \beta p \in R$. Moreover

$$\alpha[0, \omega] + \beta[\omega, 0] \in \mathfrak{Z}_{js} + (a - 1)\widehat{\mathfrak{Z}}_{js}$$

if and only if $\alpha \in R$ and $\beta \in t^{-j}R$. Using properties of the 1-cocycle $\mathfrak{f}$ it is easy to show that the two 1-cocycles $\mathfrak{f}_j$ ($j = 1, 2$):

$$\mathfrak{f}_1(a) = \alpha_1[0, \omega] + \beta_1[\omega, 0] + \mathfrak{Z}_{js}, \quad \mathfrak{f}_2(a) = \alpha_2[0, \omega] + \beta_2[\omega, 0] + \mathfrak{Z}_{js}$$

are cohomologous if and only if

$$p\alpha_1 \equiv p\alpha_2 \pmod{t^d} \quad \text{and} \quad p\beta_1 \equiv p\beta_2 \pmod{t^{d-j}},$$

where $\alpha_j, \beta_j \in F$, $p\alpha_j, p\beta_j \in R$. Note that $p = t^d\theta$.

It follows that the map $\mathfrak{f} \mapsto (p\alpha + t^dR, p\beta + t^{d-j}R)$ gives the isomorphism

$$H^1(G, \widehat{\mathfrak{Z}}_{js}) \cong R/(t^dR) \oplus R/(t^{d-j}R).$$

Therefore, according to (ii) of Lemma 1, the RG -modules $\mathfrak{Z}_{js}$ ($1 \leq j < d$) are pairwise non-isomorphic. $\square$

Now, using the description of 1-cocycles it is easy to prove the following

Corollary 3. *Put*

$$K_{\alpha, \beta}(G, \mathfrak{Z}_{js}) = \left\langle \begin{pmatrix} e & m \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} a & \alpha t^{-d}[0, \omega] + \beta t^{-d}[\omega, 0] \\ 0 & 1 \end{pmatrix} \mid m \in Z_{js} \right\rangle,$$

where α and β independently run over the representative elements of the cosets $R/(t^dR)$ and $R/(t^{d-j}R)$, respectively. Up to equivalence, the groups $K_{\alpha, \beta}(G, \mathfrak{Z}_{js})$ give all extensions of the additive group of the RG -module $\mathfrak{Z}_{js}$ by the group G .

If R is the quadratic extension of the ring of p -adic integers, then the R -representations of G were described by P.M. Gudivok (see [7]).

Finally, we have the following result

Theorem 4. *Let $\Phi_p(x)$ be decomposable into the product of at least two irreducible polynomials over R . Then the dimensions of the non-split indecomposable groups $\mathfrak{C}\mathfrak{r}\mathfrak{h}\mathfrak{s}(G, M, \mathfrak{f})$ are unbounded.*

Proof. Let $\Phi_p(x) = \eta_1(x) \cdots \eta_k(x)$ ($k > 2$) be a decomposition into a product of polynomials irreducible over R and suppose that

$$\eta_1(x) = x^n - \alpha_{n-1}x^{n-1} - \cdots - \alpha_1x - \alpha_0 \in R[x].$$

Note that $\deg(\eta_1(x)) = \deg(\eta_2(x)) = \cdots = \deg(\eta_k(x)) = n$ and $kn = p - 1$.

We will use the technic of the integral representation of finite groups, which was developed by S.D. Berman and P.M. Gudivok in [2, 3, 7].

Let ε be a primitive p th root of unity such that $\eta_1(\varepsilon) = 0$ and let r_j be a natural number, such that $\varepsilon_j = \varepsilon^{r_j}$ is a root of the polynomial $\eta_j(x)$, where $r_1 = 1$ and

$j = 1, \dots, k$. Let $\tilde{\varepsilon} = \begin{pmatrix} 0 & \cdots & 0 & \alpha_0 \\ 1 & \cdots & 0 & \alpha_1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 1 & \alpha_{n-1} \end{pmatrix}$ be the comparing matrix of $\eta_1(x)$.

The following R -representations of $G = \langle a \mid a^p = 1 \rangle$ are irreducible:

$$\delta_0 : a \mapsto 1; \quad \delta_1 : a \mapsto \tilde{\varepsilon}; \quad \delta_j : a \mapsto \tilde{\varepsilon}_j = \tilde{\varepsilon}_j^{r_j}, \quad (j = 2, \dots, k).$$

Note that the module which affords representation δ_1 is $R[\varepsilon]$ with R -basis $1, \varepsilon, \dots, \varepsilon^{n-1}$.

Let $m \in \mathbb{N}$. Define the following R -representation of $G = \langle a \rangle$ of degree $(3n+1)m$:

$$\Gamma_m : a \mapsto \begin{pmatrix} \Delta_{1m}(a) & U_m(a) \\ 0 & \Delta_{2m}(a) \end{pmatrix},$$

where

$$\Delta_{1m}(a) = \delta_0^{(m)}(a) + \delta_1^{(m)}(a) = \begin{pmatrix} E_m \otimes \delta_0(a) & 0 \\ 0 & E_m \otimes \delta_1(a) \end{pmatrix};$$

$$\Delta_{2m}(a) = \delta_2^{(m)}(a) + \delta_3^{(m)}(a) = \begin{pmatrix} E_m \otimes \delta_2(a) & 0 \\ 0 & E_m \otimes \delta_3(a) \end{pmatrix};$$

$$U_m(a) = \begin{pmatrix} E_m \otimes u & J_m(1) \otimes u \\ E_m \otimes \bar{u} & E_m \otimes \bar{u} \end{pmatrix};$$

$u = (0, 0, \dots, 0, 1)$ defines a nonzero element of $\text{Ext}(\delta_0, \delta_j)$;

$J_m(\lambda)$ is a Jordan block of degree m with λ in the main diagonal;

$\bar{u}$ is a matrix in which the first row is $(0, \dots, 0, 1)$ and all other rows are zero. The matrix $\bar{u}$ defines a nonzero element of the group $\text{Ext}(\delta_1, \delta_j)$, where $j = 2, 3$;

E_m is the unity matrix of degree m .

Lemma 5. (see [2,3]) Γ_m is an indecomposable R -representation of G .

Let $\mathfrak{W}_m = R^l$ be a module of l -dimension vectors over R affording the R -representation Γ_m . Put $\widehat{F} = F^+/R$, $\widehat{\mathfrak{W}}_m = F\mathfrak{W}_m^+/\mathfrak{W}_m$. Clearly $\widehat{F}^l \cong \widehat{\mathfrak{W}}_m$. Define $\tau : F \rightarrow F^n$ by

$$\tau(w) = w(\alpha_0, \alpha_0 + \alpha_1, \alpha_0 + \alpha_1 + \alpha_2, \dots, \alpha_0 + \dots + \alpha_{n-2}, 1), \quad (7)$$

where the α_j are coefficients of $\eta_1(x)$ and $w \in F$.

Lemma 6. (i) Each 1-cocycle of $G = \langle a \mid a^p = 1 \rangle$ at $\widehat{\mathfrak{W}}_m$ is cohomologous to a cocycle $\mathfrak{f}$, such that

$$\mathfrak{f}(a) = (X, 0, \dots, 0) + \mathfrak{W}_m,$$

where $X \in F^m$ and $pX = 0$ in $\widehat{F}^m$ (i.e. $pX \in R^m$).

(ii) Let $z \in F^n$ such that $(\tilde{\varepsilon} - E_n)z = 0$ in $\widehat{F}^n$. Then $z = \tau(w) \pmod{R^n}$, with $w \in F$ such that $\eta_1(1)w = 0$ in $\widehat{F}$.

(iii) If $V = R/(\frac{p}{\eta(1)}R)$ is the residual of ring R by the ideal $(\frac{p}{\eta(1)}R)$, then $H^1(G, \widehat{\mathfrak{W}}_n) \cong V^m$.

Proof. (i) follows from (ii) of Lemma 1. (ii) is easy to check.

(iii) by (i) we can put $\mathfrak{f}(a) = (X, 0, 0, 0)$ and $\mathfrak{g}(a) = (Y, 0, 0, 0)$, where $X = (x_1, \dots, x_n)$, $Y = (y_1, \dots, y_n)$ and $pX = pY = 0$. Note that all the equalities considered here are understood modulo the group R . Suppose that these 1-cocycles are cohomologous and $Z \in F^l$ is such that

$$(\Gamma_m(a) - E_l)Z + \mathfrak{f}(a) = \mathfrak{g}(a). \quad (8)$$

Put $Z = (Z_1, Z_2, Z_3, Z_4)$, where $Z_1 \in F^m$ and Z_2, Z_3, Z_4 are m -dimensional vectors, with i -components belong to F^n , and denoted by Z_2^i, Z_3^i and Z_4^i , respectively. By (8) we get

$$(E_m \otimes u)Z_3 + (J_m \otimes u)Z_4 + X = Y, \quad (9)$$

$$(E_m \otimes (\tilde{\varepsilon} - E_n))Z_2 + (E_m \otimes \bar{u})(Z_3 + Z_4) = 0, \quad (10)$$

$$(E_m \otimes (\tilde{\varepsilon}_2 - E_n))Z_3 = 0, \quad (E_m \otimes (\tilde{\varepsilon}_3 - E_n))Z_4 = 0. \quad (11)$$

From (11) and by (ii) we have

$$Z_3 = (\tau(v_1), \dots, \tau(v_m)), \quad Z_4 = (\tau(u_1), \dots, \tau(u_m)), \quad (12)$$

where $u_j, v_j \in F$, τ is from (7) and

$$\eta_1(1)u_j = \eta_1(1)v_j = 0. \quad (13)$$

Clearly, the equality (10) consists of m matrix equalities of the form

$$(\tilde{\varepsilon} - E_n)Z_2^i + \bar{u}\tau(w) = 0, \quad (14)$$

where $Z_2^i \in F^n$ is the i th component of Z_2 , $i = 1, \dots, m$ and $w \in F$. Since $u\tau(w) = w$ and $\bar{u}\tau(w) = (w, 0, \dots, 0)$, when all the rows of (14) are added together we obtain

$$-\eta(1)Z_2^n + w = 0, \quad (15)$$

where Z_2^n is the last component of the vector Z_2 . According to (12) and (15), (10) gives the equalities

$$-\eta(1)z_j + v_j + u_j = 0, \quad (j = 1, \dots, m) \quad (16)$$

where z_j are some components of Z_2 . From (9)

$$\begin{aligned} v_j + u_j + u_{j+1} + x_j &= y_j, & (j = 1, \dots, m-1) \\ v_m + u_m + x_m &= y_m, \end{aligned} \quad (17)$$

where $X = (x_1, \dots, x_n)$, $Y = (y_1, \dots, y_n)$ and $pX = pY = 0$. Multiplying (17) by $\eta_1(1)$ and using (16) we obtain for the components of X and Y

$$\eta_1(1)x_j = \eta_1(1)y_j \quad (j = 1, \dots, m). \quad (18)$$

Therefore, if the 1-cocycles $\mathfrak{f}$ and $\mathfrak{g}$ are cohomologous then (18) holds.

Conversely, suppose that (18) holds. Then it is not difficult to construct vectors Z_2, Z_3, Z_4 that satisfy (9) and (10), which is equivalent to (8), i.e. the 1-cocycles $\mathfrak{f}$ and $\mathfrak{g}$ are cohomologous. It follows that by going from a cocycle to an element of the cohomology group, we need to change each component in X by $\beta = \alpha \cdot p^{-1}$ modulo the group R , where $\alpha \in R$. Moreover, if $\eta_1 \cdot \beta \in R$, then must change β to 0. $\square$

Theorem 5. *Let $\varepsilon \in R$, where $\varepsilon^p = 1$ and $p > 2$. Then the description of the non-split indecomposable groups $\mathfrak{Ctns}(G, M, \mathfrak{f})$ is a wild type problem.*

Proof. For arbitrary matrices $A, B \in M(m, R)$ the map

$$\Gamma_{A,B} : a \mapsto \begin{pmatrix} E & 0 & E & A & E \\ & \varepsilon E & E & E & B \\ & & \varepsilon^2 E & 0 & 0 \\ & & & \varepsilon^3 E & 0 \\ & & & & \varepsilon^4 E \end{pmatrix}$$

is an R -representation of G of degree $l = 5m$. The R -representations $\Gamma_{A,B}$ and Γ_{A_1,B_1} are R -equivalent if and only if

$$C^{-1}AC \equiv A_1 \pmod{(1-\varepsilon)}, \quad C^{-1}BC \equiv B_1 \pmod{(1-\varepsilon)}$$

for some invertible matrix C . It follows that the description of the R -representations $\Gamma_{A,B}$ of G is a wild type problem.

For the module affording the representation $\Gamma_{A,B}$ of G we put R^l . Let X be an m -dimensional vector over F with $pX \in R^m$. Then there is a 1-cocycle $\mathfrak{f}_X : G \rightarrow \widehat{R}^l$, such that $\mathfrak{f}_X(a) = (X, 0, \dots, 0) + R^l$. The 1-cocycles $\mathfrak{f}_X$ and $\mathfrak{f}_Y$ are cohomologous if and only if

$$(1-\varepsilon)(X-Y) \in R^m.$$

Putting $X = (p^{-1}, 0, \dots, 0)$ we obtain that $H^1(G, \widehat{R}^l) \neq 0$. $\square$

REFERENCES

1. Benson, D. J., *Representations and cohomology. Cohomology of groups and modules*, Cambridge Studies in Advanced Math., 31. Cambridge University Press, Cambridge, 1998.
2. Berman, S.D., *Representations of finite groups over an arbitrary field and over rings of integers.*, Izv. Akad. Nauk SSSR Ser. Mat. **30** (1966), 69–132. (Russian)
3. Berman, S.D.; Gudivok, P.M., *Indecomposable representation of finite group over the ring p -adic integers (Russian)*, Izvestia AN USSR **28(4)** (1964), 875–910.
4. Bovdi, A.A., *Group rings (Russian)*, Kiev UMK BO, 1988, pp. 156.
5. Bovdi, V.A., Gudivok, P.M., Rudko, V.P., *Torsion free groups with indecomposable holonomy group I*, J. Group Theory **5** (2002), 75–96.
6. Bovdi, V.A., Gudivok, P.M., Rudko, V.P., *Torsion-free crystallographic groups with indecomposable holonomy group.II*, J. Group Theory **7(4)** (2004), 555–569.
7. Gudivok, P.M., *Representation of finite groups over quadratic fields*, Doklady AN USSR **5** (2002), 75–96.
8. Kopcha, G.M., *Non-split extension of the indecomposable module of p -integer representation of the cyclic group of order p^2 (Ukrainian)*, Uzhgorod University Scientific Herald. Math. Ser. **5** (2000), 49–56.

V.A. Bovdi

Institute of Mathematics, University of Debrecen

P.O. Box 12, H-4010 Debrecen, Hungary

Institute of Mathematics and Informatics, College of Nyíregyháza

Sóstói út 31/b, H-4410 Nyíregyháza, Hungary

E-MAIL: VBOVDI@MATH.KLTE.HU

V.P. Rudko

DEPARTMENT OF ALGEBRA, UNIVERSITY OF UZHGOROD

88 000, UZHGOROD, UKRAINE

E-MAIL: MATH1@UNIV.UZHGOROD.UA